

Abstract

The effects of free radicals and active oxygen species on biological molecules have created very problems in recent years. These compounds are known to be responsible for oxidative deterioration of vital molecules such as proteins, lipids and nucleic acids. Antioxidants protect biological systems against these factors and have been suggested to play a preventing role in human health. In the present study, the antioxidant activity of water and alcoholic extracts of *L.amiaceas* was determined. The extracts obtained by ultrasonic and traditional methods. 2,2-Diphenyl-1-picrylhydrazyl (DPPH) was added to the extracts and the antioxidant activity was determined by amount of reagent consumption. For tracking the progress of the reaction, high performance liquid chromatography (HPLC) was used. In order to optimize the extraction conditions, effective parameters such as extracting solvent, temperature and time were investigated. Finally, the results were analyzed by SPSS software. Analysis of variance (ANOVA) suggests that there are significant differences between extraction methods ($P < 0.05$).

Keywords: Antioxidant activity, *L.amiaceas*, Ultrasonic assisted extraction, DPPH.



University of Zabol
Graduate school
Faculty of Science
Department of Chemistry

Sc.The Thesis Submitted for the Degree of M
In the field of Analytical Chemistry

Ultra Sound Assisted Extraction and Evaluation of Antioxidant
Activity of L.amiaceas Using DPPH

Supervisor:
Dr. M. Nejati-Yazdinezhad

Advisor:
Dr. G. R. Motalleb

By:
Safiye Yosefi

March 2014